

BEST AVAILABLE COPY

AN ENANTIOSELECTIVE APPROACH TO CARBAPENEM ANTIBIOTICS: FORMAL SYNTHESIS OF (+)-THIENAMYCIN

David J. Hart* and Deok-Chan Ha

Department of Chemistry, The Ohio State University, Columbus, Ohio 43210


Summary: An enantioselective synthesis of intermediates in syntheses of thienamycin (15) and epithienamycin-C (16) is described.

Several years ago we reported that *N*-trimethylsilyl imines react with ester enolates to

as the ester component in this reaction, resulting in a direct route to 3-(1-hydroxyethyl)-2-azetidinones.² This letter presents our initial efforts to apply these observations to the area of carbapenem synthesis. Specifically, an efficient enantioselective route to β -lactam 10 and its conversion to the known thienamycin intermediate 14 is described.³⁻⁵

Ethyl (S)-8-hydroxybutyrate⁶ was converted to the corresponding dianion (2 equiv LDA, THF, -70°C) and treated with imine 1.^{7,8} The resulting crude mixture of β -lactams 2 and 3 was treated with *tert*-butyldimethylsilyl chloride and triethylamine in *N,N*-dimethylformamide⁹ to afford 4 (16%) and 5 (27%) after separation by column chromatography. Ozonolysis of 4 gave 6 (85%, mp 131-132°C) and subsequent Jones oxidation afforded acid 8 (86%, mp 105-106°C). Similar oxidative treatment of 5 gave 7 (91%) and 9 (96%, mp 146-147°C). Treatment of either 8 or 9 with lead tetraacetate¹⁰ gave 10 (mp 79-80°C) in 81% and 80% yields, respectively. From an operational standpoint it was most convenient to subject a purified mixture of β -lactams 4 and 5 to the oxidation sequence without purification of the intermediates. In this way, 10 could be prepared from 4 + 5 in a 60% overall yield.


We have demonstrated that 10 will serve as an intermediate in a synthesis of the carbapenem antibiotic thienamycin (15). Thus, treatment of 10 with *p*-nitrobenzyl 2-diazo-3-(*t*-butyldimethylsiloxy)-3-butenate and ZnCl₂ in dichloromethane gave diazoketone 11 (83%) which was converted to alcohol 12 [95%, mp 150°C (d)] upon exposure to methanol-aqueous hydrochloric acid.^{10,12} Inversion of the C-8 stereochemistry was accomplished using the excellent method of Mitsunobu.^{13,14} The resulting formate 13 (95%) was converted to the known thienamycin intermediate 14 [87%, mp 151-152°C (lit.¹² 150-152°C)] upon treatment with excess 1,2-ethanedithiol and a catalytic amount of *p*-toluenesulfonic acid in dichloromethane-isopropanol (25°C, 96h), thus completing a formal total synthesis of thienamycin (15).^{14,15} The use of 12 in a synthesis of the related carbapenem antibiotic epithienamycin-C (MM22381, 16) will appear in our full account of this work.




(a) CH₃CH(OLi)CH=C(OEt)(OLi), THF, -70°C → 25°C, 20h (b) tBuMe₂SiCl, Et₃N, DMF

(c) O₃, CH₂Cl₂, -78°C then Me₂S (d) Jones Reagent (e) Pb(OAc)₄, AcOH

We have also examined the behavior of the dianion of ethyl β-hydroxybutyrate with N-p-methoxyphenylaldimines. A few of our results are reported here. Treatment of N-p-methoxyphenylcinnamaldimine with the dianion of ethyl β-hydroxybutyrate in tetrahydrofuran (-70°C, then 20h at room temperature) afforded a mixture of 17 (25%, mp 160-161°C) and 18 + 19 (38%, mp 160-161°C, respectively).¹⁶ Silylation⁹ of the mixture of 18 + 19 gave 21 (72%, mp 97-98°C) and 22 (23%, mp 124-125°C) after separation by chromatography. The stereochemical assignment for 21 was established by sequential silylation to give 20 (88%), ceric ammonium nitrate oxidation,¹⁷ and silylation to give 5 (78%). The stereochemical assignments for the trans β-lactams were confirmed by converting both 20 and 21 to 24. Thus, ozonolysis of 21 gave 21 (86%, mp 113-115°C) and ozonolysis of 20 (90%) followed by epimerization (DBU, CH₂Cl₂, 25°C).



BEST AVAILABLE COPY



Cl, Et₃N, DMF. It is notable that we have not been able to isomerize 7 to 6 under identical conditions.

In summary, we have shown that the ester-imine condensation can be included in the list of enantioselective routes to carbapenem intermediates. The methods and intermediates described herein should also be useful in the synthesis of analogs such as carbacephems.

Acknowledgment. We thank the National Institutes of Health (AI-21074) for their support of this research. We thank Mr. Richard Weisenberger for recording mass spectra at The Ohio State University Chemical Instrument Center.

References

- Alfred P. Sloan Foundation Fellow, 1983-1987.
- Hart, D. J.; Kanai, K.; Thomas, D. T.; Yang, T.-K. *J. Org. Chem.* **1983**, *48*, 289.
- Ha, D.-C.; Hart, D. J.; Yang, T.-K. *J. Am. Chem. Soc.* **1984**, *106*, 4819.
- Preliminary accounts of this work were presented at the 18th Midwest Regional ACS Meeting in Lawrence, KS, November 2-4, 1983, Paper No. 662 and the 188th National ACS Meeting in Philadelphia, PA, August 26-31, 1984, Paper No. 115.
- Recent applications of N-silylimine chemistry to the synthesis of carbapenem intermediates prompt the disclosure of our efforts at this time: Chiba, T.; Nagatsuma, M.; Nakai, T. *Chem. Lett.* **1984**, 1927. Cainelli, G.; Contento, M.; Giacomini, D.; Panunzio, M. *Tetrahedron Lett.* **1985**, 937. Colvin, E. W.; McGarry, D. G. *J. Chem. Soc., Chem. Commun.* **1985**, 539.
- For related uses of ethyl β-hydroxybutyrate in β-lactam synthesis see Georg, G. *Tetrahedron Lett.* **1984**, 3779. Iimori, T.; Shibasaki, M. *Tetrahedron Lett.* **1985**, 1523.

6. Lemieux, R. U.; Giguere, J. *Can. J. Chem.* 1951, 29, 678. Seebach, D.; Sutter, M. A.; Weber, R. H.; Zuger, M. F. *Organic Syntheses* 1984, 63, 1. The material used in these studies was prepared by baker's yeast reduction of ethyl acetoacetate and had 86% ee. Optical rotations of intermediates will be presented in our full paper.
7. The imine was generated from cinnamaldehyde and lithium hexamethyldisilazide as described in reference 1, and used without purification.
8. For other uses of this dianion see Kraus, G. A.; Taschner, M. J. *Tetrahedron Lett.* 1977, 4575. Frater, G. *Helv. Chim. Acta* 1979, 62, 2825. Seebach, D.; Wasmuth, D. *Helv. Chim. Acta* 1980, 63, 197.
9. Corey, E. J.; Venkateswarlu, A. *J. Am. Chem. Soc.*, 1972, 94, 6190.
10. For the development of this degradation see Reider, P. U.; Schenker, L. U. *Tetrahedron Lett.* 1982, 2293.
11. For an overview of structures and syntheses see Ratcliffe, R. W.; Albers-Schönberg, G. in "Chemistry and Biology of β -Lactam Antibiotics"; Morin, R. B.; Gorman, M. Eds.; Academic Press: New York, 1982; Vol. 2, pp 227-313.
12. Ueda, Y.; Roberge, G.; Vinet, V. *Can. J. Chem.* 1984, 62, 2936.
13. Mitsunobu, O.; Kimura, J.; Iizumi, K.; Yanagida, N. *Bull. Chem. Soc. Jap.* 1976, 49, 510.
14. Melillo, D. G.; Shinkai, I.; Liu, T.; Ryan, K.; Slettinger, M. *Tetrahedron Lett.* 1980, 2783.
15. A variety of more classical acidic and basic formate hydrolysis conditions met with failure. Hydrazinolysis of 13 also gave 14 in lower yields.
16. We have observed stereochemical differences in reactions between ester enolates and N-aryl or N-silyl imines on several occasions. The mechanistic differences between these systems are under investigation.
17. Kronenthal, D. R.; Han, C. Y.; Taylor, M. K. *J. Org. Chem.* 1982, 47, 2765.
18. Aldehyde 23 was also isolated in 9% yield.

(Received in USA 8 July 1985)

BEST AVAILABLE COPY

ADSL...
than does i
ones while

During
model study
zation of t
we were unc
thermodynam
7-lactone a
mixture con

Bromol
presumably t
explored the
lactonizatio
indicated th
halogen.⁵ s
bromine and
of the other
tive formati
that the reg
group which
from the hyd

CH₃
HO